

Full Length Research Paper

Growing the threatened seahorse *Hippocampus erectus* Perry 1810 in the laboratory

Rosana Beatriz Silveira¹ and José Rodrigo Santos Silva^{1,2}

¹Laboratório de Aquicultura Marinha-LABAQUAC/Projeto Hippocampus, Porto de Galinhas, Ipojuca-PE, Brazil.

²Departamento de Estatística e Ciências Atuariais, Universidade Federal de Sergipe, São Cristóvão-SE, Brazil.

Accepted 04 June, 2016

In Brazil, a seahorse pair of *Hippocampus erectus* was collected, and in captivity, mating and copulation occurred, followed by a pregnancy period of 12 days. The fry at birth had a mean height of 7.07 ± 0.947 mm and weight of 0.0037 ± 0.0049 g. The initial feed consisted of zooplankton up to the 10th day, and subsequently, they were fed nauplii and juveniles of *Artemiasalina* and marine shrimp *Litopenaeusvannamei*. At 53 and 91 days of life, the animals had a mean height of 4.36 ± 0.315 and 6.03 ± 0.409 cm and mean weight of 0.377 ± 0.074 and 0.910 ± 0.172 g, respectively. At formation of the male's brood pouch, at 104 days of life, the mean height was 6.26 ± 0.374 cm and weight 0.9363 ± 0.1681 g. It was estimated that between 53 and 91 days of age, for every centimeter the seahorse grew, it gained about 0.3215 grams, where the equation height-weight obtained was $W=0.00731 L^{2.67728}$. Sexual maturity was reached at a height of 8.05 ± 0.56 cm. The size of the gestational period was 17.2 ± 6.70 days, and number of offspring produced was 377 ± 35.36 /pregnancy.

Keywords: Diet, height-weight ratio, fertility, brood pouch, sex ratio, *Artemiasalina*, *Litopenaeusvannamei*, Syngnathidae.

INTRODUCTION

In 1996, almost all species of seahorses (Syngnathidae: *Hippocampus*) were listed as "vulnerable" by the IUCN (International Union for Conservation of Nature), including the two Brazilian species recognized at the time: *Hippocampus reidi* and *H. erectus* (Figueiredo and Menezes, 1980; Rosa et al., 2002; Dias et al., 2002; Silveira, 2011). Currently, three Brazilian seahorse species are recognized: *H. reidi*, *H. erectus* and

H. patagonicus (Silveira et al., 2014). This inclusion in the IUCN Red List, was due to excessive commercial fishing, degradation of their environment and the ignorance of their biology in natural or captive environments (Dias et al., 2002; Baum et al., 2003; Scales, 2010; Silveira et al., 2016). Seahorses are fish threatened with extinction world wide and are among the most traded marine ornamental fishes, both as pets and for medicines and curios. However, like most marine ornamental fish, they are caught in the wild for trade because the demand is higher than the aquaculture industry can currently offer.

Global trade handles over 20 million seahorses per year, for the most varied purposes (Vincent, 1996), having reached in 2001 up to 67 tons per year, more than 23 million seahorses per year (Vincent et al., 2011). Although they are distributed all over tropical and the temperate regions worldwide (Lourie et al., 1999), 55 currently validated species (Froese and Pauly, 2016) are highly exploited, and only about 13 of the mare raised for trade (Koldewey and Martin-Smith, 2010). In 2004, CITES (International Convention on Trade in Endangered Species of Wild Fauna and Flora) included all species of seahorses in Appendix II of the Convention. The criteria of this Appendix (II) are not as restrictive as those of Appendix I, which prohibits fishing, but allows the sustainable use of resources. This decision put seahorses among the commercial marine teleosts most important to CITES (McPherson and Vincent, 2004). The purpose of CITES is to ensure that international trade does not exert unsustainable pressure on natural populations. Brazil is one of the 160 signatory countries of the Convention and is the major exporter of seahorses in Latin America, although the magnitude of this trade is not known in our country. According to 29 fishermen interviewed in Brazil, 25 said that the populations of seahorses declined around 86% (Vincent et al., 2011), but Brazilian data on these fishes are poorly known. For seahorses, there is no control or recording of commercial fishing in Brazil, with export data being insufficient to estimate the current catch of these fishes, and there is no specific protection law (Silveira, 2005). Rosa et al., (2006), working with the aquarist trade, determined the capture of 9,793 specimens of seahorses between 1997 and 2005, just in the state of Bahia, Brazil. In Pernambuco, Brazil, between 1999 and July 2000, 4,114 seahorses were exported by four companies registered with the Brazilian Institute of Environment and Renewable Natural Resources–IBAMA (Silveira, 2005). Recently, the three Brazilian species of seahorse were included in the Brazilian List of Endangered Species in the category "vulnerable" (MMA-Ordinance No. 445, 2014).

The trade of these fish is very lucrative for middlemen, who pay very little to the humble fishermen who live off or supplement their family income with this fishing. A live seahorse is sold in ornamental fish stores for about R\$ 20.00, while the fisherman is paid R\$ 1.00 for the "darker" and R\$ 4.00 for the "colored", preferably red (Silveira, 2005). Fishing is done on any age or size, with no sparing of pregnant males. In open markets, for example in Recife, or Paráor Baía deParanaguá, PR, one can find the trade of specimens of the three species occurring in Brazil, sold as dehydrated parts for use in "folk medicine," curios, lucky charms, etc., with price ranging between R\$ 4.00 and R \$10.00 per unit, depending on the size and skeletal condition (Silveira 2005, Silveira et al., 2014). In addition to the trade in ornamental fish and

medicinal products, the animals suffer from environmental degradation and fishing by catch, especially with trawling (Salin et al., 2005, Foster and Vincent 2012; Vincent et al., 2011; Jardim et al., 2012, Filizand Taskavak 2012, Foster and Arreguin-Sánchez 2013). To reduce the pressure on wild populations of seahorses, it has been proposed to breed them for trade (Job et al., 2002; Koldewey and Martin-Smith, 2010).

Hippocampus erectus, despite its taxonomic uncertainty and distribution (Boehm et al., 2013; Silveira, 2011; Silveira et al., 2014) and all other threats common to the Syngnathidae and several other groups, such as trawling (Baum et al., 2003; Foster and Vincent, 2012), its culture has been demonstrated to be adequate and easy to manage to meet commercial and conservation needs (Lin et al., 2009). In the coastal waters of Pernambuco, Brazil, the species is sighted with great difficulty, and over the past 13 years of research, our team found only six live specimens and three dead (one on the beach and two in trawls). Of the live specimens, only one was male. In this paper, we described the biology of *H. erectus* and its first rearing in captivity in Brazil.

METHODS

Collection, food and acclimatization of seahorses

Only one pair of seahorses of the species *H. erectus* was found and collected on the beach of Porto de Galinhas, Ipojuca, Pernambuco. The animals were not found together, they were caught one month apart. The seahorses underwent prophylactic treatment with 0.1% copper sulfate (Duijn, 1956) in individual quarantines and fed 10- to 14-day post-larval marine shrimp *Litopenaeus vannamei* (PLs 10-14) and adult *Artemiasalina*. Afterwards, the seahorses were kept in a 60-l aquarium containing natural sea water sterilized by chlorination, with salinity of 30, pH 8.3, photoperiod of 9h, biological filter plate and moderate aeration, in a room at 25°C ± 1. The levels of ammonia, nitrite and nitrate were monitored weekly with colorimetric kits (Alcon, Brazil), and kept at 0 mg/l. After six months of acclimatization in the laboratory, the male courted the female, resulting in mating between the pair.

Estimate of the gestation period and average brood produced

The development of six pregnancies of the same pair was monitored to determine the size of the gestational period and the average number fry expelled from the brood pouch at end of pregnancy.

Aquariums for newborns and juveniles

The birth of offspring, 100 fingerlings were transferred to

four 100-liters (25 animals per tank, 0.25 ind/l), with external filter built with skeleton of calcareous algae *Halimeda* sp and parts of ceramic such as filter elements, and addition of skimmer and 11-W UV filter, with recirculating natural seawater. The water parameters were monitored every three days and maintained as with breeding tanks. In the water recirculation, 10% of the water was exchanged every other day until the animals were 3 months old, when they were transferred to two 500-liters fitted as the previous ones. At this stage, a *Halimeda* sp background was added and the tanks were planted with green seaweed. There are no more water changes, only fresh water replenishment for salinity maintenance.

Feeding newborns and juveniles

The initial feeding of newborns consisted of wild zooplankton collected in the sea and at low tide with 150- μ m mesh net. The samples were collected, homogenized and diluted daily to 120 ± 20 ind/ml (mostly cope pods and copepodites), providing 400 ml of this dilution, twice daily (10:00 and 16:00h). The photoperiod of the newborns was increased to 12h, so that they could spend more time feeding. The seahorses consumed only wild zooplankton up to the 10th day of life, and from the 11th day and henceforth, nauplii of *A. salina* and *L. vannamei* (50% of each crustacean) were introduced at a concentration of 80 ± 20 ind/ml, with 500 ml of this mixture being offered twice daily (10:00 and 16:00h). Feeding with zooplankton was continued for seven days to minimize the effect of diet change. After the second month of life, the seahorses were fed 5- to 7-day post-larval marine shrimp (PLs) and *A. salina* juveniles (0.5 cm long), and starting on the fourth month of life, 10-day PLs and *A. salina* adults.

Biometrics and weight

The offspring were weighed and measured at birth (n=40) and at 53 (n=50) and 91 (n=84) days of life. The height (Ht), linear measurement from the top of the head to the tip of the stretched tail (Lourie et al., 1999) was made with a plastic ruler, which was submerged to avoid removing the animal out of the water. Weighing was done with a digital semi-analytical balance (0.001g), on which the animal was placed after the removal of excess water with tissue paper. Newborns, on the other hand, were weighed on an analytical balance (0.0001 g).

Height determination on brood pouch formation

The animals used for the calculation of mean height at brood pouch formation were juveniles from the smallest seahorse with fully formed pouch to the largest with the pouch still in formation. Juveniles with small pouch

formation (values below the smallest male with fully formed pouch) were not considered for calculation because we wanted to determine the mean height of the seahorses with a fully formed pouch (Figure 1).

Statistical Analysis

The mean height at brood pouch formation was estimated by linear regression, using the various classes in height, and the variable indicating brood pouch formation assumed two values, "1" for pouch formed and "0" for pouch in formation.

The equation for the weight/height ratio was obtained by linear regression, which estimated the variation in weight according to height of seahorses, using a 5% significance level. The software used was R3.2.2 (The R Core Team 2015).

RESULTS

Courting and copulation by *H. erectus* occurred during the day, where the male courted the female with characteristic moves, which were similar to those observed in this laboratory for *H. reidi*. Copulation occurred with the transfer of hydrated oocytes into the male's brood pouch, where they were fertilized. According to six consecutive pregnancies, the mean time for the species was 17.2 ± 6.70 days, and 377 ± 35.36 offspring resulted in each pregnancy. The fry were expelled through the orifice of the brood pouch after various and sequential contractions. The new born shadan mean height of 7.02 ± 0.43 mm and all exhibited a homogeneous light yellow color. They were fed from the first moment with wild zoo plankton without any difficulty and no deaths occurred. As of the 11th day, nauplii of *L. vannamei* and *A. salina* were introduced as food, which were well accepted by the seahorses. From day 11 today 18, 16 animals died and were the only deaths recorded in three of the four experimental aquariums. On completing 53 days of life, the fish had a mean height of 4.36 ± 0.31 cm and wet weight of 0.377 ± 0.074 g (Fig. 2). The first brood pouch sign in males was recorded after 74 days when there was a different color in the first ring of the ventral surface of the tail. After 91 days, the animals exhibited a height of 6.03 ± 0.409 cm and weighed 0.910 ± 0.172 g, but no pouch was fully formed, which occurred at 107 days. The formation of the male brood pouch occurred at a mean height of 6.26 cm, and weight of 0.9363 ± 0.1681 g. Data analysis demonstrated that 95% of males of this species begin brood pouch formation between 5.90 and 6.10 cm, while the fully formed pouch appears between 6.024 and 6.498 (Table 1). The sex ratio of offspring was 1:1, where 40 males and 44 females were observed ($Z = 0.4364$, $p = 0.6631$).




Figure 1. Illustrative scheme of the criteria used in selecting the data to estimate the average height brood pouch formation of males: [] brood pouch in formation; [] smaller male with brood pouch fully formed; [] brood pouch fully formed; [] larger male with brood pouch in formation; the numbers represent the heights (cm).




Figure 2. Height (a) and wet weight (b) of the seahorse *Hippocampus erectus* cultivated from birth to 91 days of life (n=84).


Considering a linear model for the weight x height equation, $W = -1.024 + 0.3215 \times H_t$, it was estimated that during the intense growth period (between 53 and 91 days of age), for every centimeter the seahorse grew, it gained about 0.321g. The general equation obtained for

the height-weight ratio, $W = 0.0067H^{2.7247}$ (adjusted $r^2 = 0.9863$, $n = 174$, $p < 2.2e^{-16}$), showed that in this phase, all juveniles during the second month of life developed dermal appendages, which persisted in adulthood but

Table 1. Height at formation of the brood pouch in the *Hippocampus erectus* male.

	Coefficient	Standard error	t	P	Lower 95%	Upper 95%
Interception*	6.017	0.056	108.419	0.000	5.906	6.129
Pouch**	0.244	0.118	2.065	0.043	0.007	0.481

*Mean height of male brood pouch in formation (cm); **Mean difference between the animals with pouch formed and those with pouch in formation (cm). Mean height of animals with fully formed pouch is given by the sum of the coefficients $6.017 + 0.244 = 6.261$



decreased with age (Fig. 4). When sexual maturity was reached at 8.05 ± 0.56 cm, courtship and copulation began between pairs, but there was a high degree of promiscuity with males courting males, even in the presence of females.

DISCUSSION

According to Foster and Vincent (2004), the majority of seahorse species exhibit monogamous behavior, but there is an apparently low level of polygamy recorded for

H. abdominalis, *H. subelongatus*, *H. fuscus* and *H. guttulatus*, and there was one polygamous event for *H. reidi* (Silveira, 2009). Courting and mating between males and females of *H. erectus* in our study followed the pattern described for the genus (Strawn, 1958; Mason Jones and Lewis, 1996; Garrick-Maidonent, 1997; Foster and Vincent, 2004; Lin et al., 2008; Silveira, 2009; Rosenqvist and Berglund, 2011). However, after the offspring reached adulthood and started reproductive activities (at a height of about 8.0 cm), there was a high level of promiscuity with males courting males, even in



Figure 4. Young of *Hippocampus erectus* cultivated for 150 days.

the presence of the same number of females (1:1). These observations were very intriguing, as it involved choices made by the males repeatedly for many days. A similar situation had only been observed for *H. reidi*, but that was induced by the unique isolation of males (Silveira, 2009). The mean height (Ht) of cultured *H. erectus* at birth has been reported to be 15.9 mm (Herald and Racowiks 1951), 16.4 ± 1.8 mm (Vite-Garcia et al., 2014), 13.0 mm standard length (SL) (Chung et al., 1989) and 11.3 ± 0.06 mm (SL) (Lin et al., 2008). In the present study, *H. erectus* had a mean height of 7.02 ± 0.43 mm at birth, smaller than that reported by the above cited authors.

According to Vincent and Giles (2003), the juvenile size in the same species can vary with latitude and environmental variables, with photoperiod and temperature also being factors, where higher latitudes result in increased size of egg and juvenile at birth. In fact, the difference in size at birth between *H. erectus* grown in Pernambuco (8°S) and other countries such as the US (23°N , Florida; 37°N , San Francisco) could be explained by this condition, and this probably would be due to the need for greater gain in energy reserves and fat for protection against the low temperatures that occur at high latitudes. Though growing up in captivity under controlled conditions, characteristics are preserved due to genetic baggage responsible for the adaptation of each

community to its environment over millions of years (Freeman and Herron, 2009).

In our work, *H. erectus* measured 4.36 ± 0.31 cm (Ht) at 53 days of life, with a mean weight of 0.377 ± 0.074 g, values similar to those of 4.34 ± 0.2 cm and 0.249 ± 0.053 g found by Vite-Garcia et al., (2014) at 60 days of life for the same species, fed on a diet *Artemia* sp. enriched with Selco[®], where in this phase, 7 days or more of culture can produce a large difference in final mean height. Herald and Racowiks (1951) found 5.08 cm for *H. erectus* at 60 days of life. According to Lin et al., (2008), *H. erectus* had a standard length (SL) of 6.32 ± 0.52 cm at 67 days and wet weight was 0.45 ± 0.12 g.

In the third month, the heights were very similar, 6.35 cm and 6.03 ± 0.401 cm for Herald and Racowiks (1951) and this work, respectively. Regarding the formation of the male brood pouch, Chung et al., (1989) observed at 10.0 cm (SL), corresponding to an age of three months, the initial formation of this structure. Herald and Racowiks (1951) reported the formation of the brood pouch between 6.35 to 7.0 cm in height, corresponding to 3.5 months, while Scarratt (1995) estimated the formation of this structure between 5 and 6 months for *H. erectus*. For *H. guttulatus*, Cartagena (2014) observed the first sign of pouch formation at 124 days of life (4 months) with 8.7 cm in height. In this work, the first sign of a brood pouch (different color on the ventral side of the first caudal rings)

appeared at 74 days of life (2.46 months), where the mean height at brood pouch formation was 6.26 cm, corresponding to 3.5 months of age. In situ data showed *H. erectus* with a formed brood pouch at 4.0 cm in height (Teixeira and Musick, 2001).

In this work, after following six consecutive pregnancies of the same parents, the mean length of gestation was determined to be 17.20 ± 6.70 days, but by observing the exact day of three clutches inside the male brood pouch (of the six recorded) until the day of birth, two showed 12 days and one 13. For the other three clutches, the exact date of the transfer of eggs was not observed. Thus, it is possible that the average gestational period was overestimated and that *H. erectus* had the same gestational period as *H. reidi* in Pernambuco (12 days) (Silveira and Fontoura, 2010). The mean number of offspring produced in the six pregnancies was 377 ± 35.36 juveniles. Lin et al., (2008) estimated for *H. erectus* (six pairs) a gestational period of 17.33 ± 2.94 and offspring of 272.33 ± 66.45 juveniles per brood pouch. Lin et al., (2008) reported for their cultivation conditions a survival rate of $71.11 \pm 10.18\%$ up to nine weeks of culture at 28°C , Vite-Garcia et al. (2014) obtained 82% survival in the culture of *H. erectus* up to 60 days of live, at $26 \pm 1^\circ\text{C}$, while in this work, we observed 84% survival at 150 days of life at $25 \pm 1^\circ\text{C}$. Hora and Joyeux (2009) used a diet consisting of wild zooplankton, *A. salina* enriched with Super Selco[®] and the mysid *Misidyum gracile* (live and frozen), and obtained 88.3% survival in *H. reidi* culture. Pham and Lin (2013) achieved a survival rate of 84.3% for *H. reidi* up to 28 days of live using a diet of *Artemia* enriched with Dan's Feed[®] and 74.5% with a diet of wild plankton, in both cases stocked at 1 ind/l. Zang et al., (2010) stocked *H. erectus* at 1 ind/L and obtained 87.8% survival up to 40 days of life.

The relationship between estimated weight and height in this work, $W = 0.0067 H^{2.7247}$ (adjusted $r^2 = 0.9863$, $n = 174$, $p < 2.2e^{-16}$) was similar to that obtained by Lin et al., (2008), $W = 0.0034 L^{2.5535}$ ($r^2 = 0.9903$, $n = 12$, $p < 0.01$), although we observed a faster weight gain in our cultures, possibly due to the quality of food provided (wild zooplankton, *L. vannamei* nauplii and post-larvae fed artificial diet, *A. salina* nauplii, juveniles and adults enriched with microalgae in this work versus *A. salina* nauplii, juveniles, adults enriched with microalgae and fish oil emulsion-Super Selco[®], and frozen *Mysis* spp. in Lin et al., 2008). Palma et al., (2012) observed better development for *H. guttulatus* fed with the shrimp *Palaemonetes varians*, with artificial diet for 12 weeks, recording the number of pregnancies, greater number of fingerlings incubated and greater size on birth also.

According to Teixeira and Musick (2001), the fertility of *H. erectus* in Chesapeake Bay, FL was 451 ± 232.10 embryos per pregnancy, with a maximum of 1552, suggesting that there is the possibility to optimize the reproductive capacity of the species in culture, by

working with selected specimens, proper nutrition and enriched diet.

We believe that the aquaculture often endangered species, especially those with high commercial appeal, such as seahorses and many other ornamental fishes and marine invertebrates, should be encouraged as a conservation measure for the species in their environment.

ACKNOWLEDGEMENTS

Thanks to Petrobras, through the Petrobras Socio ambiental Program. Thanks to Chico Mendes Institute of Biodiversity Conservation for research license No. 40311-1. Dr. A. Leyva helped with English translation and editing in the preparation of the manuscript.

REFERENCES

- Baum JK, Meeuwig JJ, Vincent ACJ (2003). Bycatch of lined seahorses (*Hippocampus erectus*) in a Gulf of Mexico shrimp trawl fishery. Fish. B-NOAA 101: 721-731.
- Boehm JT, Woodall L, Teske PR, Lourie SA, Baldwin C, Waldman J, Hickerson M (2013). Marine dispersal and barriers drive Atlantic seahorse diversification. J. Biogeogr. 40: 1-11.
- Cartagena AB (2014). Rearing of the seahorse *Hippocampus guttulatus*: Key factors involved in growth and survival. Research thesis, Universitat de les Illes Balears: 292p.
- Chung KS, Correa M, Manrique R (1989). Cultivo experimental del caballito de mar *Hippocampus erectus*. Bol. Inst. Oceanogr. Venezuela. 28: 191-731.
- Dias TL, Rosa IL, Baum JK (2002) Threatened Fishes of the World: *Hippocampus erectus* Perry, 1810 (Syngnathidae). Environ. Biol. Fishes 65:326.
- Duijn CV (1956). Diseases of Fishes, Water Life, London.
- Figueiredo JL, Menezes NA (1980). Manual de peixes marinhos do sudeste do Brasil. III. Teleostei (2), Museu de Zoologia da Universidade São Paulo, São Paulo. 90 pp.
- Filiz H, Taskavak E (2012). Field surveys on recent situation of seahorses in Turkey. Biharean Biologist, 6 (1): 55-60.
- Foster SJ, Vincent ACJ (2004). Life history and ecology of seahorses: implications for conservation and management. J. Fish. Biol. 65: 1-61.
- Foster SJ, Vincent ACJ (2012). Advice in spite of great uncertainty: assessing and addressing bycatch of small fishes with limited data using *Stellifer illecebrosus* as a case study. Aquatic Conserv: Mar. Freshw. Ecosyst. 22: 639-651.
- Foster SJ, Arreguin-Sánchez F (2013). Using distribution patterns of small fishes to assess small fish by-catch in

- tropical shrimp trawl fisheries. *Animal Conservation*: 1-8.
- Freeman S, Herron JC (2009). *Análise Evolutiva*. 4. Ed. Porto Alegre: Artmed, pp 848.
- Froese, R. and D. Pauly (eds.) (2016). *Fish Base*. World Wide Web electronic publication. Version 01/2016. Accessed at www.fishbase.org, 20 April 2016.
- Garrick-Maidonent N (1997). *Seahorses: conservation and care*. United Kingdom.
- Herald E, Rakowicz M (1951). Stable requirements for raising sea horses. *Aquarium J*. 22: 234-242.
- Hora MDSC Da, Joyeux, JC (2009). Closing the reproductive cycle: Growth of the seahorse *Hippocampus reidi* (Teleostei, Syngnathidae) from birth to adulthood under experimental conditions. *Aquaculture* 292: 37-41.
- IUCN (2015). The IUCN Red List of threatened species. Version 2015-4. International Union for Conservation of Nature. Accessed at <http://www.iucnredlist.org>, 19 May 2016.
- Jardim E, Prista N, Fernandes AC, Silva D, Ferreira A L, Abreu P, Fernandes P (2012). Manual de procedimentos a bordo: arrasto de fundo com portas. *Relat. Cient.Téc. Inst. Invest. Pescas Mar Série digital* (<http://inrb.pt/ipimar>), nº 55:20 p. + Anexos.
- Job SD, Do HH, Meeuwig JJ, Hall HJ (2002). Culturing the oceanic seahorse, *Hippocampus kuda*. *Aquaculture* 214: 333-341.
- Koldewey HJ, Martin-Smith KM (2010). A global review of seahorse aquaculture. *Aquaculture* 302, 131-152.
- Lin Q, Lin J, Zhang D (2008). Breeding and juvenile culture of the lined seahorse, *Hippocampus erectus* Perry, 1810. *Aquaculture* 277: 287-292.
- Lin Q, Zhang D, Lin J (2009). Effects of light intensity, stocking density, feeding frequency and salinity on the growth of sub-adult seahorses *Hippocampus erectus* Perry, 1810. *Aquaculture* 292: 111-116.
- Lourie S, Vincent ACJ, Hall HJ (1999). *Seahorses: an identification guide to the world's species and their conservation*, Project Seahorse, London.
- McPherson JM, Vincent ACJ (2004). Assessing East African trade in seahorses species as a basis for conservation under international controls. *Aquatic Conserv: Mar. Freshw. Ecosyst*. 14: 521-538.
- Masonjones HD, Lewis SM (1996). Courtship Behavior in the Dwarf Seahorse, *Hippocampus zosterae*. *Copeia* 3: 634- 640.
- MMA-Ministério do Meio Ambiente-(2014). Fauna brasileira ameaçada de extinção. Anexos à Portaria 445 do Ministério do Meio Ambiente, de 17/12/2004, D.O.U. nº 245, Seção I, pág. 126, de 18/12/2014.
- Palma, J., Andrade, J. P. and Bureau, DP (2012). Growth, Reproductive Performances, and Brood Quality of Long Snout Seahorse, *Hippocampus guttulatus*, Fed Enriched Shrimp Diets. *J. World Aquacult. Soc.* 43: 802–813.
- Pham NK, Lin J (2013). The Effects of Different Feed Enrichments on Survivorship and Growth of Early Juvenile Longsnout Seahorse, *Hippocampus reidi*. *J. World Aquacult. Soc.* 44 (3): 435-446.
- Rosa IL, Dias T, Baum J (2002). Threatened fishes of the world: *Hippocampus reidi* Ginsburg, 1933 (Syngnathidae). *Environ. Biol. Fishes* 64, 378.
- Rosa IL, Sampaio CLS, Barros AT (2006). Collaborative monitoring of the ornamental trade of seahorses and pipefishes (Teleostei: Syngnathidae) in Brazil: Bahia State as a case study. *Neotrop.Ichthyol.* 4: 247-252.
- Rosenqvist G, Berglund A (2011). Sexual signals and mating patterns in Syngnathidae. *J. Fish. Biol.* 78: 1647–1661.
- Salin KR, Yohannan TM, Nair CM (2005). Fisheries and trade of seahorses, *Hippocampus* spp., in southern India. *Fish. Manag. Ecol.* 12: 269–273
- Scales, H (2010). Advances in the ecology, biogeography and conservation of seahorses (genus *Hippocampus*). *ProgPhysGeog.* 34(4): 443–458.
- Scarratt AM (1995) Techniques for raising lined seahorse (*Hippocampus erectus*). *Aqr. Front.* 3: 24-29.
- Silveira RB (2005). Dinâmica populacional do cavalo-marinho *Hippocampus reidi* (Syngnathidae) no manguezal de Maracaípe, Ipojuca, PE. In: Departamento de Zoologia. Pontifícia Universidade Católica do Rio Grande do Sul, Porto Alegre.
- Silveira RB (2009). Sobre o comportamento sexual do cavalo-marinho *Hippocampus reidi* Ginsburg 1933 (Pisces:Syngnathidae). *Biociências* 17: 20-32.
- Silveira RB (2011). Registros de cavalos-marinhos (Syngnathidae: *Hippocampus*) ao longo da costa brasileira. *Oecol. Aust.* 15 (2): 316-325.
- Silveira RB, Fontoura NF (2010). Fecundity and fertility of the longsnout seahorse, *Hippocampus reidi* (Teleostei: Syngnathidae), in tropical Brazil. *R. bras. Bioci.* 8 (4): 362-367.
- Silveira RB, Siccha-Ramirez R, Silva JRS, Oliveira C (2014). Morphological and molecular evidence for the occurrence of three *Hippocampus* species (Teleostei: Syngnathidae) in Brazil. *Zootaxa* 3861: 317-332.
- Silveira RB, Silva JRS, Fontoura, NF (2016). Reproductive period, average height for the development of the brood pouche and sexual maturation of the seahorse *Hippocampus reidi* (Syngnathidae) in the Northeast of Brazil. *AARJMD* 3 (3): 140-155.
- Strawn K (1958). Life history of the pigmy *Hippocampus zosterae* Jordan and Gilbert at Cedar Key Florida. *Copeia* 16-22.
- Teixeira RL, Musick JA (2001). Reproduction and food habits of the lined seahorse, *Hippocampus erectus* (Teleostei: Syngnathidae) of Chesapeake Bay, Virginia. *Bras J. Biol.* 61: 79-90.
- The R Core Team (2015). *R: A Language and Environment*

- for Statistical Computing, R Foundation for Statistical Computing, Vienna, Austria.
- Vincent ACJ (1996). The International Trade in Seahorses, TRAFFIC International.
- Vincent ACJ, Foster SJ, Koldewey HJ (2011). Conservation and management of seahorses and other Syngnathidae. *J. Fish. Biol.* 78: 1681-1724.
- Vincent ACJ, Giles BG (2003). Correlates of reproductive success in a wild population of *Hippocampus whitei*. *J. Fish. Biol.* 63: 344-355.
- Vite-Garcia N, Simoes N, Arjona O, Mascaro M, Palacios E (2014). Growth and survival of *Hippocampus erectus* (Perry, 1810) juveniles fed on *Artemia* with different HUFA levels. *Lat. Am. J. Aquat. Res.* 42 (1): 150-159.
- Zhang, D, Zhang Y, Lin J, Lin Q(2010). Growth and survival of juvenile lined seahorse, *Hippocampus erectus* (Perry), at different stocking densities. *Aquacult. Res.* 42:9–13.